

Boost your performance and confidence with these topic-based exam questions

Practice questions created by actual examiners and assessment experts

Detailed mark scheme

Suitable for all boards

Designed to test your ability and thoroughly prepare you

Level: IGCSE Oxford AQA Biology (9201)

Subject: Biology

Topic: IGCSE AQA Biology



To be used by all students preparing for IGCSE Oxford AQA Biology (9201)
Students of other Boards may also find this useful

Biology

IGCSE AQA

Key skills



1 1	d absorb oxygen through their skin.	
(a) What is the name of th	e process in which oxygen enters the skin cells?	
Tick one box.		
Active transport		
Diffusion		
Osmosis		
Respiration		
		1)
The table below shows informati	ion about four skin cells of an earthworm.	

Cell	Percentage of oxygen			
Cell	Outside cell	Inside cell		
Α	9	8		
В	12	8		
С	12	10		
D	8	12		

PRACTICE

(b) Which cell has the smallest difference in percentage of oxygen between the outside and the inside of the cell?

Tick one box.

АВ	С	D
----	---	---

(1)



1.	Earthworms are small animals that live in soil. Earthworms have no specialised exchange system and absorb oxygen through their skin.	l gas
(c)	Which cell will oxygen move into the fastest?	
	Tick one box.	
	A B C D	
		(1)
(d)	Earthworms have a large surface area to volume ratio.	
	Suggest why a large surface area to volume ratio is an advantage to an earthworm.	
		(1)
(e)	The earthworm uses enzymes to digest dead plants.	. ,
(6)		
	Many plants contain fats or oils.	
	Which type of enzyme would digest fats?	
	Copyright © 1984 Exam Papers Macrice	
		(1)
		. ,



Earthworms are small animals that live in soil. Earthworms have no specialised gas

L.	e	exchange system and absorb oxygen through their skin.	
(f)	Earth	nworms move through the soil.	
	This	movement brings air into the soil.	
		d plants decay faster in soil containing earthworms compared with soil containing no nworms.	
	Expla	ain why.	
			(3)
	E	XAM PAPERS PRACTICE	
(g)	When	earthworms reproduce, a sperm cell from one earthworm fuses with an	
egg c	ell fron	n a different earthworm.	
	Nam	e the process when an egg cell and a sperm cell fuse.	
			- (1)
	(h)	Some types of worm reproduce by a process called fragmentation.	
		In fragmentation, the worm separates into two or more parts. Each part grows into a new worm.	
		What type of reproduction is fragmentation?	
			(1) arks)



- **1.** Earthworms are small animals that live in soil. Earthworms have no specialised gas exchange system and absorb oxygen through their skin.
- **2.** A student carried out an investigation using chicken eggs.

This is the method used.

- 1. Place 5 eggs in acid for 24 hours to dissolve the egg shell.
- 2. Measure and record the mass of each egg.
- 3. Place each egg into a separate beaker containing 200 cm³ of distilled water.
- 4. After 20 minutes, remove the eggs from the beakers and dry them gently with a paper towel.
- 5. Measure and record the mass of each egg.

Table 1 shows the results.

Table 1

	Egg	Mass of egg without shell in grams	Mass of egg after 20 minutes in grams	
	1	73.5	77.0	
ľ	2	70.3	73.9	IC
	3	72.4	75.7	
	4	71.6	73.1	
	5	70.5	73.8	

(a) Another student suggested that the result for egg **4** was anomalous.

Do you agree with the student?

Give a reason for your answer



culate the percenta	ige change in mass o	of egg 3.		
	Percentage chan	ge in mass =		
plain why the mass	es of the eggs increa	sed.		
XAM	PAPE	RS P	RAC1	FICE
e nyh 1984 I Sain Papers Practice				



- **1.** Earthworms are small animals that live in soil. Earthworms have no specialised gas exchange system and absorb oxygen through their skin.
- (d) Explain how the student could modify the investigation to determine the concentration of the solution inside each egg.

EXAM PAPERS PRACTICE

Chicken egg shells contain calcium. Calcium ions are moved from the shell into the cytoplasm of the egg.

Table 2 shows information about the concentration of calcium ions.

Table 2

Location	Concentration of calcium ions in arbitrary units
Egg shell	0.6
Egg cytoplasm	2.1

(3)



(e)		how calcium ions are moved from the shell into the cytoplasm of the egg.	-
			-
			-
		(та	- (3 otal 12 marks
3.		dent investigated the effect of different sugar solutions on potato tissue.	
	1.	Add 30 cm ³ of 0.8 mol dm ⁻³ sugar solution to a boiling tube.	
	2.	Repeat step 1 with equal volumes of 0.6, 0.4 and 0.2 mol dm ⁻³ sugar solut	ions.
	3.	Use water to give a concentration of 0.0 mol dm ⁻³ .	
	4.	Cut five cylinders of potato of equal size using a cork borer.	
	5.	Weigh each potato cylinder and place one in each tube.	
	6.	Remove the potato cylinders from the solutions after 24 hours.	
	7.	Dry each potato cylinder with a paper towel.	
	8.	Reweigh the potato cylinders.	

The table below shows the results.



1. Earthworms are small animals that live in soil. Earthworms have no specialised gas exchange system and absorb oxygen through their skin.

Concentration of sugar solution in mol dm ⁻³	Starting mass in g	Final mass in g	Change of mass in g	Percentage (%) change
0.0	1.30	1.51	0.21	16.2
0.2	1.35	1.50	0.15	х
0.4	1.30	1.35	0.05	3.8
0.6	1.34	1.28	-0.06	-4.5
0.8	1.22	1.11	-0.11	-9.0

	_
	_
Percentage change in mass = %	

(b) Why did the student calculate the percentage change in mass as well as the change in grams?




1.

Earthworms are small animals that live in soil. Earthworms have no specialised gas exchange system and absorb oxygen through their skin.

(1)

- (c) Complete the graph using data from the table above.
 - Choose a suitable scale and label for the x-axis.
 - Plot the percentage (%) change in mass.



(4)



1.	 Earthworms are small animals that live in soil. Earthworms have no speci exchange system and absorb oxygen through their skin. 	alised gas
(d)	Use your graph to estimate the concentration of the solution inside the potato cells.	
	Concentration = mol dm ⁻³	(1)
		(-7
(e)	The results in the table above show the percentage change in mass of the	
potato	o cylinders.	
	Explain why the percentage change results are positive and negative.	
	EXAM PAPERS PRACTIC	E
	Cup 1611 © 2024 Com Papers Practice	
		(3)
(f)	Suggest two possible sources of error in the method given above.	
	1	-
		_
	2	-
		_
	(то	(2) otal 13 marks)



- **1.** Earthworms are small animals that live in soil. Earthworms have no specialised gas exchange system and absorb oxygen through their skin.
- **4.** Cells, tissues and organs are adapted to take in different substances and get rid of different substances.

The table shows the concentration of four ions outside cells and inside cells.

Ion	Concentration outside cells in mmol per dm ³	Concentration inside cells in mmol per dm ³
Sodium	140	9
Potassium	7	138
Calcium	2	27
Chloride	118	3

(a) Use information from the table above to complete the following sentences.

Sodium ions will move into cells by the process

of PAPERS	PRACTICE
Potassium ions will move into cells by the process	
of	

(b) Some students investigated the effect of the different concentrations of sugar in four drinks, **A**, **B**, **C** and **D**, on the movement of water across a partially permeable membrane.

The students:

- made four bags from artificial partially permeable membrane
- put equal volumes of 5% sugar solution in each bag
- weighed each bag containing the sugar solution
- placed one bag in each of the drinks, A, B, C and D
- after 20 minutes removed the bags containing the sugar solution and weighed them again.

For more help visit our website https://www.exampaperspractice.co.uk/

(2)



1.	exchange system and absorb oxygen through their skin.	as
Th	ne diagram below shows how they set up the investigation.	
Artific perm bag, ends	cial partially neable membrane tied at both Sugar solution A 0.1% sugar solution A 0.1% sugar solution A 0.1% sugar solution Solution Solution A 0.1% sugar solution Solution Solution Solution	
(i)	The bag in drink A got heavier after 20 minutes.	
	Explain why.	
	EXAM PAPERS PRACTICE	
(ii)	In which drink, A , B , C or D , would you expect the bag to show the smallest change in mass? Tick (✓) one box.	(3)
	A B C D	

(1)



Diagra	m A shows the	e same cell of a concell in a hypotonic same cell in a hypotonic	solution.		 (2) (Total 8 marks)
Diagra	m A shows the	cell in a hypotonic same cell in a hype	solution.		
Diagra	m B shows the	same cell in a hype	ertonic solution.		
Diagra EX					
EX	Diagram A	ADEL			
e seri (un		APER	RS PR	iagram B	ICE
E S	SURFORMATION OF THE PARTY OF TH	1000			
a) What is a	hypertonic solu	tion?			




Earthworms are small animals that live in soil. Earthworms have no specialised gas

(i)	a hypotonic solution
(ii)	a hypertonic solution?
plain	what has happened to the plant cell in diagram B.
plain	what has happened to the plant cell in diagram B.
plain	what has happened to the plant cell in diagram B.
xplain	what has happened to the plant cell in diagram B.



- **1.** Earthworms are small animals that live in soil. Earthworms have no specialised gas exchange system and absorb oxygen through their skin.
- (d) Animal cells will also change when placed in different solutions.

Some red blood cells are put in a hypotonic solution.



(4)

(Total 12 marks)

For more help visit our website https://www.exampaperspractice.co.uk/