



EXAM PAPERS PRACTICE

Boost your performance and confidence with these topic-based exam questions

Practice questions created by actual examiners and assessment experts

Detailed mark scheme

Suitable for all boards

Designed to test your ability and thoroughly prepare you

Level: HL IB in Biology

Subject: Biology

Topic: IB HL Biology

Type: Mark Scheme

2002

XVIII

1583

All International Baccalaureate IB Topic Questions HL Biology

BIOLOGY

HL - IB

Key skills

Answer 1

The correct answer is B

Load is the weight of the person acting downwards, the fulcrum is the pivot point around the ball of the foot and the effort is the contracting force from the muscle.

Answer 2

The correct answer is D

D is not a movement demonstrated by simple synovial joints such as the knee or elbow as inversions are complex movements made by joints in the foot.

Answer 3


The correct answer is D

The distance between the two Z lines is the same as the length of a sarcomere; when the muscle contracts, this length decreases as the actin slides across the myosin filaments to increase the overlap between the two fibres. This reduces the visible size of the light band. The length of the dark band does not change because it spans the length of the myosin filaments which do not change in length when contraction occurs.

Answer 4

One myosin powerstroke requires hydrolysis of one molecule of ATP and moves one actin filament 20 nm.

How many ATP molecules would be required to move a single actin filament 0.3 μm when the sarcomere contracts?

- 
- 1000nm in 1 μm
1. Convert 0.3 μm into nm $\rightarrow 0.3 \times 1000 = 300\text{nm}$
 2. Calculate how many powerstrokes have occurred to move the actin 300nm $\rightarrow 300 \div 20 = 15$ powerstrokes
 3. Multiply by the number of ATP produced per power stroke $\rightarrow 15 \times 1 = 15$ ATP

The correct answer is B

Copyright © 2024 Exam Papers Practice

Answer 5

The correct answer is D

Although A mentions active transport (which does require ATP) it is not correct because calcium ions diffuse in through voltage gated channels.

Calcium is responsible for exposing the myosin binding sites (as seen in the last question).

Crossbridge formation occurs as a result of the exposure of binding sites caused by calcium.

Answer 6

The correct answer is B

Don't be put off by having not heard of phosphocreatine before; the key to answering this is in the question thread. Phosphocreatine provides a source of phosphate. This means that the increased force of contraction shown in the graph can be attributed to a higher availability of phosphate to make ATP for muscle contraction: $ADP + P_i = ATP$. The information is not enough to say whether mice run faster as a result of the phosphocreatine (A) and answers C and D are incorrect.

Answer 7

The correct answer is A

The transverse section shows the two different fibres, actin and myosin, represented by the thick and thin dots and A is in the dark band of the micrograph, where the myosin and actin overlap.

B is a light band made up of actin only.

C is in the very centre of the sarcomere; this region will contain only myosin filaments.

D is the same as B.